

Controlled G-Frames and Their G-Multipliers in Hilbert spaces

Asghar Rahimi^a, Abolhassan Fereydooni^b

^aDepartment of Mathematics

University of Maragheh

Maragheh, Iran

rahimi@maragheh.ac.ir,

^bDepartment of Mathematics

Vali-e-Asr University of Rafsanjan

Rafsanjan, Iran

Fereydooniman@yahoo.com

Abstract

Multipliers have been recently introduced by P. Balazs as operators for Bessel sequences and frames in Hilbert spaces. These are operators that combine (frame-like) analysis, a multiplication with a fixed sequence (called the symbol) and synthesis. Weighted and controlled frames have been introduced to improve the numerical efficiency of iterative algorithms for inverting the frame operator. Also g -frames are the most popular generalization of frames that include almost all of the frame extensions.

In this manuscript the concept of the controlled g -frames will be defined and we will show that controlled g -frames are equivalent to g -frames and so the controlled operators C and C' can be used as preconditions in applications. Also the multiplier operator for this family of operators will be introduced and some of its properties will be shown.

1 Introduction

In [31], R. Schatten provided a detailed study of ideals of compact operators by using their singular decomposition. He investigated the operators

⁰**2000 Mathematics Subject Classification:** Primary: 42C15; Secondary: 41A58, 47A58.

Keywords: frame; g -frame; g -Bessel; g -Riesz basis; g -orthonormal basis; multiplier; Schatten p -class; Hilbert-Schmidt; trace class; controlled frames; weighted frame; controlled g -frame, (C, C') -controlled g -frame, (C, C') -controlled g -multiplier operator.

of the form $\sum_k \lambda_k \varphi_k \otimes \overline{\psi_k}$ where (ϕ_k) and (ψ_k) are orthonormal families. In [3] the orthonormal families were replaced with Bessel and frame sequences to define Bessel and frame multipliers.

Definition 1.1 *Let \mathcal{H}_1 and \mathcal{H}_2 be Hilbert spaces, let $(\psi_k) \subseteq \mathcal{H}_1$ and $(\phi_k) \subseteq \mathcal{H}_2$ be Bessel sequences. Fix $m = (m_k) \in l^\infty$. The operator $\mathbf{M}_{m,(\phi_k),(\psi_k)} : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ defined by*

$$\mathbf{M}_{m,(\phi_k),(\psi_k)}(f) = \sum_k m_k \langle f, \psi_k \rangle \phi_k$$

*is called the **Bessel multiplier** for the Bessel sequences (ψ_k) and (ϕ_k) . The sequence m is called the symbol of \mathbf{M} .*

Several basic properties of these operators were investigated in [3]. Multipliers are not only interesting from a theoretical point of view, see e.g. [3, 18, 13], but they are also used in applications, in particular in the field of audio and acoustic. They have been investigated for fusion frames [1], for generalized frames [29], p -frames in Banach spaces [30] and for Banach frames [16, 17]. In signal processing they are used for Gabor frames under the name of Gabor filters [23], in computational auditory scene analysis they are known by the name of time-frequency masks [24]. In real-time implementation of filtering system they approximate time-invariant filters [7]. As a particular way to implement time-variant filters they are used for example for sound morphing [12] or psychoacoustical modeling [8].

G -frames, introduced by W. Sun in [32] and improved by the first author [28], are a natural generalization of frames which cover many other extensions of frames, e.g. bounded quasi-projectors [19, 20], pseudo-frames [22], frame of subspaces or fusion frames [10], outer frames [2], oblique frames [11, 15], and a class of time-frequency localization operators [14]. Also it was shown that g -frames are equivalent to stable spaces splitting studied in [27]. All of these concepts are proved to be useful in many applications. Multipliers for g -frames introduced in [29] and some of its properties investigated.

Weighted and controlled frames have been introduced recently to improve the numerical efficiency of iterative algorithms for inverting the frame operator on abstract Hilbert spaces [6], however they are used earlier in [9] for spherical wavelets. In this manuscript the concept of controlled g -frame will be defined and we will show that any controlled g -frame is equivalent a g -frame and the role of controller operators are like the role of preconditions matrices or operators in linear algebra. Furthermore the multiplier operator for these family will be investigated.

The paper is organized as follows. In Section 2 we fix the notation in this paper, summarize known and prove some new results needed for the rest of the paper. In Section 3 we will define the concept of controlled g -frames and we will show that a controlled g -frame is equivalent to a g -frame and so the controlling operators can be used as precondition matrices in the problems related to applications. In section 4 we will define multipliers of controlled g -frame operators and we will prove some of its properties.

2 Preliminaries

Now we state some notations and theorems which are used in the present paper. Through this paper, \mathcal{H} and \mathcal{K} are Hilbert spaces and $\{\mathcal{H}_i : i \in I\}$ is a sequence of Hilbert spaces, where I is a subset of \mathbb{Z} . $\mathcal{L}(\mathcal{H}, \mathcal{K})$ and $\mathcal{L}(\mathcal{H})$ is the collection of all bounded linear operators from \mathcal{H} into \mathcal{K} and \mathcal{H} respectively.

A bounded operator T is called *positive* (respectively *non-negative*), if $\langle Tf, f \rangle > 0$ for all $f \neq 0$ (respectively $\langle Tf, f \rangle \geq 0$ for all f). Every non-negative operator is clearly self-adjoint. For $T_1, T_2 \in \mathcal{L}(\mathcal{H})$, we write $T_1 \leq T_2$ whenever

$$\langle T_1(f), f \rangle \leq \langle T_2(f), f \rangle, \quad \forall f \in \mathcal{H}.$$

If $U \in \mathcal{L}(\mathcal{H})$ is non-negative, then there exists a unique non-negative operator V such that $V^2 = U$. Furthermore V commutes with every operator that commute with U . This will be denoted by $V = U^{\frac{1}{2}}$. Let $\mathcal{GL}(\mathcal{H})$ be the set of all bounded operators with a bounded inverse and $\mathcal{GL}^+(\mathcal{H})$ be the set of positive operators in $\mathcal{GL}(\mathcal{H})$. For $U \in \mathcal{L}(\mathcal{H})$, $U \in \mathcal{GL}^+(\mathcal{H})$ if and only if there exists $0 < m \leq M < \infty$ such that

$$m \leq U \leq M.$$

For U^{-1} we have

$$M^{-1} \leq U^{-1} \leq m^{-1}.$$

The following theorem can be found in [21].

Theorem 2.1 *Let $T_1, T_2, T_3 \in \mathcal{L}(\mathcal{H})$ and $T_1 \leq T_2$. Suppose $T_3 > 0$ commutes with T_1 and T_2 then*

$$T_1 T_3 \leq T_2 T_3.$$

Recall that if T is a compact operator on a separable Hilbert space \mathcal{H} , then in [31] it is proved that there exist orthonormal sets $\{e_n\}$ and $\{\sigma_n\}$ in \mathcal{H} such that

$$Tx = \sum_n \lambda_n \langle x, e_n \rangle \sigma_n,$$

for $x \in \mathcal{H}$, where λ_n is the n -th singular value of T . Given $0 < p < \infty$, the **Schatten p -class** of \mathcal{H} [31], denoted \mathcal{S}_p , is the space of all compact operators T on \mathcal{H} with the singular value sequence $\{\lambda_n\}$ belonging to ℓ^p . It was shown that [33], \mathcal{S}_p is a Banach space with the norm

$$\|T\|_p = \left[\sum_n |\lambda_n|^p \right]^{\frac{1}{p}}. \quad (1)$$

\mathcal{S}_1 is called the *trace class* of \mathcal{H} and \mathcal{S}_2 is called the *Hilbert-Schmidt class*. $T \in \mathcal{S}_p$ if and only if $T^p \in \mathcal{S}_1$. Moreover $\|T\|_p^p = \|T^p\|_1$. Also, $T \in \mathcal{S}_p$ if and only if $|T|^p = (T^*T)^{\frac{p}{2}} \in \mathcal{S}_1$ if and only if $T^*T \in \mathcal{S}_{\frac{2}{p}}$. Moreover, $\|T\|_p^p = \|T^*\|_p^p = \| |T| \|_p^p = \| |T|^p \|_1 = \|T^*T\|_{\frac{2}{p}}$.

2.1 G -Frames

For any sequence $\{\mathcal{H}_i : i \in I\}$, we can assume that there exists a Hilbert space \mathcal{K} such that for all $i \in I$, $\mathcal{H}_i \subseteq \mathcal{K}$ (for example $\mathcal{K} = (\bigoplus_{i \in I} \mathcal{H}_i)_{\ell^2}$).

Definition 2.1 A sequence $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ is called *generalized frame*, or simply a **g -frame**, for \mathcal{H} with respect to $\{\mathcal{H}_i : i \in I\}$ if there exist constants $A > 0$ and $B < \infty$ such that

$$A\|f\|^2 \leq \sum_{i \in I} \|\Lambda_i f\|^2 \leq B\|f\|^2, \quad \forall f \in \mathcal{H}. \quad (2)$$

The numbers A and B are called *g -frame bounds*.

$\Lambda = \{\Lambda_i : i \in I\}$ is called *tight g -frame* if $A = B$ and *Parseval g -frame* if $A = B = 1$. If the second inequality in (2) holds, the sequence is called *g -Bessel sequence*.

$\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ is called a **g -frame sequence**, if it is a g -frame for $\overline{\text{span}}\{\Lambda_i^*(\mathcal{H}_i)\}_{i \in I}$.

It is easy to see that, if $\{f_i\}_{i \in I}$ is a frame for \mathcal{H} with bounds A and B , then by putting $\mathcal{H}_i = \mathbb{C}$ and $\Lambda_i(\cdot) = \langle \cdot, f_i \rangle$, the family $\{\Lambda_i : i \in I\}$ is a g -frame for \mathcal{H} with bounds A and B . Let

$$\left(\bigoplus_{i \in I} \mathcal{H}_i \right)_{\ell_2} = \left\{ \{f_i\}_{i \in I} \mid f_i \in \mathcal{H}_i, \forall i \in I \text{ and } \sum_{i \in I} \|f_i\|^2 < +\infty \right\}. \quad (3)$$

Proposition 2.2 [25] $\Lambda = \{ \Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I \}$ is a g -Bessel sequence for \mathcal{H} with bound B , if and only if the operator

$$T_\Lambda : \left(\bigoplus_{i \in I} \mathcal{H}_i \right)_{\ell_2} \longrightarrow \mathcal{H}$$

defined by

$$T_\Lambda(\{f_i\}_{i \in I}) = \sum_{i \in I} \Lambda_i^*(f_i)$$

is a well-defined and bounded operator with $\|T_\Lambda\| \leq \sqrt{B}$. Furthermore

$$T_\Lambda^* : \mathcal{H} \longrightarrow \left(\bigoplus_{i \in I} \mathcal{H}_i \right)_{\ell_2}$$

$$T_\Lambda^*(f) = \{\Lambda_i f\}_{i \in I}.$$

If $\Lambda = \{ \Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I \}$ is a g -frame, the operators T_Λ and T_Λ^* in Proposition 2.2 are called **synthesis operator** and **analysis operator** of $\Lambda = \{ \Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I \}$, respectively.

Proposition 2.3 [26] $\Lambda = \{ \Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I \}$ is a g -frame for \mathcal{H} if and only if the synthesis operator T_Λ is well-defined, bounded and onto.

The ordinary version of the next theorem which is proved in [16], can be extended easily to the general case.

Theorem 2.2 [16] $\Lambda = \{ \Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I \}$ is a g -Bessel sequence for \mathcal{H} if and only if the operator

$$S_\Lambda : \mathcal{H} \longrightarrow \mathcal{H}, \quad S_\Lambda = \sum_{i \in I} \Lambda_i^* \Lambda_i f, \quad (4)$$

is a welldefined operator. In this case S_Λ is bounded.

Theorem 2.3 [17] $\Lambda = \{ \Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I \}$ is a g -frame for \mathcal{H} if and only if the operator

$$S_\Lambda : \mathcal{H} \longrightarrow \mathcal{H}, \quad S_\Lambda = \sum_{i \in I} \Lambda_i^* \Lambda_i f,$$

is a welldefined invertible operator. In this case S_Λ is bounded.

S_Λ is called the **g -frame operator** of $\Lambda = \{\Lambda_i : i \in I\}$ and it is known [25] that S_Λ is a positive and

$$AI \leq S_\Lambda \leq BI,$$

where A and B are the frame bounds. Every $f \in \mathcal{H}$ has an expansion $f = \sum_i \Lambda_i^* \Lambda_i S_\Lambda^{-1} f$. One of the most important advantages of g -frames is a resolution of identity $\sum_i \Lambda_i^* \Lambda_i S_\Lambda^{-1} = I$.

2.2 Multipliers of g -frames

The concept of multipliers for g -Bessel sequences introduced by the first author in [29] and some of their properties will be shown.

Definition 2.4 Let $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ and $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be g -Bessel sequences. If for $m \subset \mathbb{C}$, the operator

$$\begin{aligned} \mathbf{M} &= \mathbf{M}_{m, \Lambda, \Theta} : \mathcal{H} \rightarrow \mathcal{H} \\ \mathbf{M}(f) &= \sum_i m_i \Lambda_i^* \Theta_i f \end{aligned} \tag{5}$$

is welldefined, then \mathbf{M} is called the **g -multiplier** of Λ, Θ and m .

If $m = (m_i) = (1, 1, 1, \dots)$ and $\mathbf{M} = I$, (Λ, Θ) is called a *pair dual* (i.e. $I = \sum_{i \in I} \Lambda_i^* \Theta_i$).

Let $\{\lambda_i\}$ and $\{\varphi_i\}$ be Bessel sequences and $m \in \ell^\infty$, consider the corresponding g -Bessel sequences $\Lambda_i \cdot = \langle \cdot, \lambda_i \rangle$ and $\Theta_i \cdot = \langle \cdot, \varphi_i \rangle$. For any $f \in \mathcal{H}$ we have:

$$\mathbf{M}_{m, \Lambda, \Theta}(f) = \mathbf{M}_{m, (\lambda_k), (\varphi_k)}(f) = \sum_i m_i \langle f, \varphi_i \rangle \lambda_i.$$

It is easy to show that the adjoint of $\mathbf{M}_{m, \Lambda, \Theta}$ is $\mathbf{M}_{\overline{m}, \Theta, \Lambda}$.

Lemma 2.5 [29] If $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ is a g -Bessel sequence with bound B_Θ and $m = (m_i) \in \ell^\infty$, then $\{m_i \Theta_i\}_{i \in I}$ is a g -Bessel sequence with bound $\|m\|_\infty B_\Theta$.

Like weighted frames [6], $\{m_i\Theta_i\}_{i \in I}$ can be called *weighted g-frame (g-Bessel)*. By using the synthesis and the analysis operators of Λ and $m\Theta$, respectively, we can write

$$\mathbf{M}_{m,\Lambda,\Theta}f = \sum_i m_i \Lambda_i^* \Theta_i f = \sum_i \Lambda_i^* (m_i \Theta_i) f = T_\Lambda \{m_i \Theta_i f\} = T_\Lambda T_{m\Theta}^* f.$$

So

$$\mathbf{M}_{m,\Lambda,\Theta} = T_\Lambda T_{m\Theta}^*. \quad (6)$$

If we define the diagonal operator

$$\begin{aligned} D_m : \left(\bigoplus \mathcal{H}_i \right)_{\ell_2} &\rightarrow \left(\bigoplus \mathcal{H}_i \right)_{\ell_2}, \\ D_m((\xi_i)) &= (m_i \xi_i)_{i \in I} \end{aligned} \quad (7)$$

then

$$\mathbf{M}_{m,\Lambda,\Theta} = T_\Lambda D_m T_\Theta^*. \quad (8)$$

The notations in (6), (7) and (8) were used for proving the following propositions in [29].

Proposition 2.6 [29] *Let $m \in \ell^\infty$, $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be a g-Riesz base and $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be a g-Bessel sequence. The map $m \rightarrow \mathbf{M}_{m,\Lambda,\Theta}$ is injective.*

Proposition 2.7 [29] *Let $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ and $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be g-Bessel sequences for \mathcal{H} . If $m = (m_i) \in c_0$ and $(\text{rank} \Theta_i) \in \ell^\infty$, then $\mathbf{M}_{m,\Lambda,\Theta}$ is compact.*

Proposition 2.8 [29] *Let $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ and $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be g-Bessel sequences for \mathcal{H} . If $m = (m_i) \in \ell^p$ and $(\dim \mathcal{H}_i)_{i \in I} \in \ell^\infty$, then $\mathbf{M}_{m,\Lambda,\Theta}$ is a Schatten p -class operator.*

Corollary 2.9 [29] *Let $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ and $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be g-Bessel sequences for \mathcal{H} .*

1. *If $m = (m_i) \in \ell^1$ and $(\dim \mathcal{H}_i)_{i \in I} \in \ell^\infty$, then $\mathbf{M}_{m,\Lambda,\Theta}$ is a trace- class operator.*
2. *If $m = (m_i) \in \ell^2$ and $(\dim \mathcal{H}_i)_{i \in I} \in \ell^\infty$, then $\mathbf{M}_{m,\Lambda,\Theta}$ is a Hilbert-Schmit operator.*

3 Controlled g -frames

Weighted and controlled frames have been introduced recently to improve the numerical efficiency of iterative algorithms for inverting the frame operator. In [6], it was shown that the controlled frames are equivalent to standard frames and it was used in the sense of preconditioning.

In this section, the concepts of controlled frames and controlled Bessel sequences will be extended to g -frames and we will show that controlled g -frames are equivalent g -frames.

Definition 3.1 *Let $C, C' \in \mathcal{GL}^+(\mathcal{H})$. The family $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ will be called a (C, C') -controlled g -frame for \mathcal{H} , if Λ is a g -Bessel sequence and there exists constants $A > 0$ and $B < \infty$ such that*

$$A\|f\|^2 \leq \sum_{i \in I} \langle \Lambda_i C f, \Lambda_i C' f \rangle \leq B\|f\|^2, \quad \forall f \in \mathcal{H}. \quad (9)$$

*A and B will be called controlled frame bounds. If $C' = I$, we call $\Lambda = \{\Lambda_i\}$ a **C -controlled g -frame** for \mathcal{H} with bounds A and B . If the second part of the above inequality holds, it will be called **(C, C') -controlled g -Bessel sequence** with bound B .*

The proof of the following lemmas is straightforward.

Lemma 3.1 *Let $C \in \mathcal{GL}^+(\mathcal{H})$. The g -Bessel sequence $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ is (C, C) -controlled Bessel sequence (or (C, C) -controlled g -frame) if and only if there exists constant $B < \infty$ (and $A > 0$) such that*

$$\sum_{i \in I} \|\Lambda_i C f\|^2 \leq B\|f\|^2, \quad \forall f \in \mathcal{H}$$

$$(\text{ or } A\|f\|^2 \leq \sum_{i \in I} \|\Lambda_i C f\|^2 \leq B\|f\|^2, \quad \forall f \in \mathcal{H}).$$

We call the (C, C) -controlled Bessel sequence and (C, C) -controlled g -frame, C^2 -controlled Bessel sequence and C^2 -controlled g -frame with bounds A, B .

Lemma 3.2 *For $C, C' \in \mathcal{GL}^+(\mathcal{H})$, the family $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ is a (C, C') -controlled Bessel sequence for \mathcal{H} if and only if the operator*

$$L_{CAC'} : \mathcal{H} \rightarrow \mathcal{H}, \quad L_{CAC'} f := \sum_{i \in I} C' \Lambda_i^* \Lambda_i C f,$$

is well defined and there exists constant $B < \infty$ such that

$$\sum_{i \in I} \langle \Lambda_i C f, \Lambda_i C' f \rangle \leq B \|f\|^2, \quad \forall f \in \mathcal{H}.$$

The operator

$$L_{C\Lambda C'} : \mathcal{H} \rightarrow \mathcal{H}, \quad L_{C\Lambda C'} f := \sum_{i \in I} C' \Lambda_i^* \Lambda_i C f,$$

is called the (C, C') -controlled Bessel sequence operator, also $L_{C\Lambda C'} = C S_{\Lambda} C'$. It follows from the definition that for a g -frame, this operator is positive and invertible and

$$A I \leq L_{C\Lambda C'} \leq B I.$$

Also, if C and C' commute with each other, then C', C'^{-1}, C, C^{-1} commute with $L_{C\Lambda C'}, C S_{\Lambda}, S_{\Lambda} C'$.

The following proposition shows that any g -frame is a controlled g -frame and versa. This is the most important advantage of weighted and controlled g -frame in the sense of precondition.

Proposition 3.3 *Let $C \in \mathcal{GL}^+(\mathcal{H})$. The family $\Lambda = \{ \Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I \}$ is a g -frame if and only if Λ is a C^2 -controlled g -frame.*

Proof. Suppose that Λ is a C^2 -controlled g -frame with bounds A, B . Then

$$A \|f\|^2 \leq \sum_{i \in I} \|\Lambda_i C f\|^2 \leq B \|f\|^2, \quad \forall f \in \mathcal{H},$$

For $f \in \mathcal{H}$

$$\begin{aligned} A \|f\|^2 &= A \|C C^{-1} f\|^2 \leq A \|C\|^2 \|C^{-1} f\|^2 \leq \|C\|^2 \sum_{i \in I} \|\Lambda_i C C^{-1} f\|^2 \\ &= \|C\|^2 \sum_{i \in I} \|\Lambda_i f\|^2. \end{aligned}$$

Hence

$$A \|C\|^{-2} \|f\|^2 \leq \sum_{i \in I} \|\Lambda_i f\|^2, \quad \forall f \in \mathcal{H}.$$

On the other hand for every $f \in \mathcal{H}$,

$$\sum_{i \in I} \|\Lambda_i f\|^2 = \sum_{i \in I} \|\Lambda_i C C^{-1} f\|^2 \leq B \|C^{-1} f\|^2 \leq B \|C^{-1}\|^2 \|f\|^2.$$

These inequalities yields that Λ is a g -frame with bounds $A\|C\|^{-2}, B\|C^{-1}\|^2$. For the converse assume that Λ is g -frame with bounds A', B' . Then for all $f \in \mathcal{H}$,

$$A'\|f\|^2 \leq \sum_{i \in I} \|\Lambda_i f\|^2 \leq B'\|f\|^2.$$

So for $f \in \mathcal{H}$,

$$\sum_{i \in I} \|\Lambda_i C f\|^2 \leq B'\|C f\|^2 \leq B'\|C\|^2\|f\|^2.$$

For lower bound, the g -frameness of Λ shows that for any $f \in \mathcal{H}$,

$$A'\|f\|^2 = A'\|C^{-1}C f\|^2 \leq A'\|C^{-1}\|^2\|C f\|^2 \leq \|C^{-1}\|^2 \sum_{i \in I} \|\Lambda_i C f\|^2.$$

Therefore Λ is a C^2 -controlled g -frame with bounds $A'\|C^{-1}\|^{-2}, B'\|C\|^2$. \square

Proposition 3.4 *Assume that $\Lambda = \{\Lambda_i : i \in I\}$ is a g -frame and $C, C' \in \mathcal{GL}^+(\mathcal{H})$, which commute with each other and commute with S_Λ . Then $\Lambda = \{\Lambda_i : i \in I\}$ is a (C, C') -controlled g -frame.*

Proof. Let Λ be g -frame with bounds A, B and $m, m' > 0, M, M' < \infty$ so that

$$m \leq C \leq M, \quad m' \leq C' \leq M'.$$

Then

$$mA \leq CS_\Lambda \leq MB,$$

because C commute with S_Λ . Again C' commutes with CS_Λ and then

$$mm'A \leq L_{C\Lambda C'} \leq MM'B.$$

\square

4 Multipliers of Controlled g -frames

Extending the concept of multipliers of frames, in this section, we will define controlled g -frame's multiplier for C -controlled g -frames in Hilbert spaces. The definition of general case (C, C') -controlled g -frames goes smooth.

Lemma 4.1 *Let $C, C' \in \mathcal{GL}^+(\mathcal{H})$ and $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$, $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be C'^2 and C^2 -controlled g -Bessel sequences for \mathcal{H} , respectively. Let $m \in \ell^\infty$. The operator*

$$\mathbf{M}_{mC\Theta\Lambda C'} : \mathcal{H} \rightarrow \mathcal{H}$$

defined by

$$\mathbf{M}_{mC\Theta\Lambda C'} f := \sum_{i \in I} m_i C \Theta_i^* \Lambda_i C' f$$

is a well-defined bounded operator.

Proof. Assume $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$, $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be C'^2 and C^2 -controlled g -Bessel sequences for \mathcal{H} with bounds B, B' , respectively. For any $f, g \in \mathcal{H}$ and finite subset $J \subset I$,

$$\begin{aligned} \left\| \sum_{i \in J} m_i C \Theta_i^* \Lambda_i C' f \right\| &= \sup_{g \in \mathcal{H}, \|g\|=1} \left\| \sum_{i \in J} m_i \langle \Lambda_i C' f, \Theta_i C^* g \rangle \right\| \\ &\leq \sup_{g \in \mathcal{H}, \|g\|=1} \sum_{i \in J} |m_i| \|\Lambda_i C' f\| \|\Theta_i C^* g\| \\ &\leq \sup_{g \in \mathcal{H}, \|g\|=1} \|m\|_\infty \left(\sum_{i \in I} \|\Theta_i C^* g\|^2 \right)^{\frac{1}{2}} \left(\sum_{i \in J} \|\Lambda_i C' f\|^2 \right)^{\frac{1}{2}} \\ &\leq \|m\|_\infty \sqrt{BB'} \|f\| \end{aligned}$$

This shows that $\mathbf{M}_{mC\Theta\Lambda C'}$ is well-defined and

$$\|\mathbf{M}_{mC\Theta\Lambda C'}\| \leq \|m\|_\infty \sqrt{BB'}.$$

□

Above Lemma ia a motivation to define the following definition.

Definition 4.2 *Let $C, C' \in \mathcal{GL}^+(\mathcal{H})$ and $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$, $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be C'^2 and C^2 -controlled g -Bessel sequences for \mathcal{H} , respectively. Let $m \in \ell^\infty$. The operator*

$$\mathbf{M}_{mC\Theta\Lambda C'} : \mathcal{H} \rightarrow \mathcal{H}$$

defined by

$$\mathbf{M}_{mC\Theta\Lambda C'} f := \sum_{i \in I} m_i C \Theta_i^* \Lambda_i C' f, \quad (10)$$

is called the (C, C') -controlled multiplier operator with symbol m .

By using representations (6) and (8), we have

$$\mathbf{M}_{mC\Theta\Lambda C'} = C\mathbf{M}_{m\Theta\Lambda}C' = CT_{\Theta}D_mT_{\Lambda}^*C'.$$

The proof of Proposition 4.7. of [29] shows that if $m = (m_i) \in \ell^p$ and $(\dim \mathcal{H}_i)_{i \in I} \in \ell^\infty$, then the diagonal operator D_m is a Schatten p -class operator. Since \mathcal{S}_p is a $*$ -ideal of $\mathcal{L}(\mathcal{H})$ so we have:

Theorem 4.3 *Let $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ and $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be controlled g -Bessel sequences for \mathcal{H} . If $m = (m_i) \in \ell^p$ and $(\dim \mathcal{H}_i)_{i \in I} \in \ell^\infty$, then $\mathbf{M}_{mC\Theta\Lambda C'}$ is a Schatten p -class operator.*

And

Corollary 4.4 *Let $\Lambda = \{\Lambda_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ and $\Theta = \{\Theta_i \in \mathcal{L}(\mathcal{H}, \mathcal{H}_i) : i \in I\}$ be controlled g -Bessel sequences for \mathcal{H} .*

1. *If $m = (m_i) \in \ell^1$ and $(\dim \mathcal{H}_i)_{i \in I} \in \ell^\infty$, then $\mathbf{M}_{mC\Theta\Lambda C'}$ is a trace-class operator.*
2. *If $m = (m_i) \in \ell^2$ and $(\dim \mathcal{H}_i)_{i \in I} \in \ell^\infty$, then $\mathbf{M}_{mC\Theta\Lambda C'}$ is a Hilbert-Schmit operator.*

Acknowledgment: Some of the results in this paper were obtained during the first author visited the Acoustics Research Institute, Austrian Academy of Sciences, Austria, he thanks this institute for their hospitality.

References

- [1] M. L. Arias and M. Pacheco, *Bessel fusion multipliers Journal of Mathematical Analysis and Applications*, 348(2), pp.581–588, 2008.
- [2] A. Aldroubi, C. Cabrelli and U. Molter, *Wavelets on irregular grids with arbitrary dilation matrices and frame atomics for $L^2(\mathbb{R}^d)$* , *Appl. Comput. Harmon. Anal.*, 17, pp. 119–140, 2004.
- [3] P. Balazs, *Basic definition and properties of Bessel multipliers* *Journal of Mathematical Analysis and Applications* 325(1), pp. 571–585, 2007.

- [4] P. Balazs, *Hilbert-Schmidt Operators and Frames-Classifications, Approximation by Multipliers and Algorithms*, *International Journal of Wavelets, Multiresolution and Information Processing*, 6(2), pp. 315–330, 2008.
- [5] P. Balazs, *Matrix Representations of Operators Using Frames. Sampling Theory in Signal and Image Processing (STSIP)*, 7(1), pp.39–54, 2008.
- [6] P. Balazs, J-P. Antoine and A. Grybos, *Weighted and Controlled Frames*. *International Journal of Wavelets, Multiresolution and Information Processing*, Volume: 8, Issue: 1, pp. 109–132, 2010.
- [7] P. Balazs, W. A. Deutsch, A. Noll, J. Rennison, and J. White, *STx Programmer Guide, Version: 3.6.2*. Acoustics Research Institute, Austrian Academy of Sciences, 2005.
- [8] P. Balazs, B. Laback, G. Eckel, and W. A. Deutsch, *Introducing time-frequency sparsity by removing perceptually irrelevant components using a simple model of simultaneous masking*. *IEEE Transactions on Audio, Speech and Language Processing*, forthcoming:–, 2009.
- [9] I. Bogdanova, P. Vandergheynst, J-P. Antoine, L. Jacques and M. Morvidone, *Stereographic wavelet frames on the sphere*, *Applied Comput. Harmon. Anal.*, 19: pp. 223–252, 2005.
- [10] P. G. Casazza and G. Kutyniok, *Frames of Subspaces,* *Wavelets, Frames and Operator Theory* (College Park, MD, 2003), pp. 87–113, *Contemp. Math.* 345, Amer. Math. Soc., Providence, RI, 2004.
- [11] O. Christensen and Y. C. Eldar, *Oblique dual frames and shift-invariant spaces*, *Appl. Comput. Harmon. Anal.* 17, pp. 48–68, 2004.
- [12] Ph. Depalle, R. Kronland-Martinet, and B. Torr sani, *Time-frequency multipliers for sound synthesis*. In *Proceedings of the Wavelet XII conference, SPIE annual Symposium, San Diego*, August 2007.
- [13] M. D rfler and B. Torresani, *Representation of operators in the time-frequency domain and generalized gabor multipliers*. *J. Fourier Anal. Appl.*, accepted, 2009.
- [14] D rfler, *Gabor analysis for a class of signals called music*, Ph.D. thesis, University of Vienna, 2003.

- [15] Y. C. Eldar, *Sampling with arbitrary sampling and reconstruction spaces and oblique dual frame vectors*, *J. Fourier Anal. Appl.* 9, pp.77–96, 2003.
- [16] A. Fereydooni, A. Safapour, Banach Pair frames, preprint.
- [17] A. Fereydooni, A. Safapour, Pair frames, preprint arXiv: 1109.3766v1.
- [18] H. G. Feichtinger and K. Nowak, *A first survey of Gabor multipliers, chapter 5, pp. 99-128*. Birkhauser, Boston, 2003.
- [19] M. Fornasier, *Decomposition of Hilbert spaces: local construction of global frames*, *Proc. Int. Conf. on Constructive function theory, Varna, B. Bojanov Ed., DARBA, Sofia*, pp. 275–281, 2003.
- [20] M. Fornasier, *Quasi-orthogonal decomposition of structured frames*, *J. Math. Anal. Appl.*, 289, pp. 180–199, 2004.
- [21] Heuser, H.: *Functional analysis*. John Wiley, New York, 1982.
- [22] S. Li and H. Ogawa, *Pseudoframes for subspaces with application*, *J. Fourier Anal. Appl.*, 10, pp. 409–431, 2004.
- [23] G. Matz and F. Hlawatsch, *Linear Time-Frequency Filters: On-line Algorithms and Applications*, chapter 6 in 'Application in Time-Frequency Signal Processing', pp. 205–271. eds. A. Papandreou-Suppappola, Boca Raton (FL): CRC Press, 2002.
- [24] G. Matz, D. Schafhuber, Karlheinz Gröchenig, M. Hartmann, and Franz Hlawatsch. Analysis, Optimization, and Implementation of Low-Interference Wireless Multicarrier Systems. *IEEE Trans. Wireless Comm.*, 6(4): pp.1–11, 2007.
- [25] A. Najati and A. Rahimi, *Generalized Frames in Hilbert spaces*, Bulletin of Iranian Mathematical Society, 35, No. 1, pp. 97–109, 2009.
- [26] A. Najati, M.H. Faroughi and A. Rahimi, *G-frames and stability of g-frames in Hilbert spaces*, *Methods of Functional Analysis and Topology*, 14, pp. 271–286, 2008.
- [27] P. Oswald, *Multilevel Finite Element Approximation: Theory and Application*, *Teubner Skripten zur Numerik*, Teubner, Stuttgart, 1994.
- [28] A. Rahimi, *Frames and Their Generalizations in Hilbert and Banach Spaces*, LAP Lambert Academic Publishing, 2011.

- [29] A. Rahimi, *Multipliers of Genralized frames in Hilbert spaces*, Bulletin of the Iranian Mathematical Society, 37 (1), pp. 63–8-, 2011.
- [30] A. Rahimi and P. Balazs, *Multipliers of p -Bessel sequences in Banach spaces*, Integral Equations and Operator Theory , Volume 68, No. 2, pp. 193-205, 2010.
- [31] R. Schatten, *Norm Ideals of Completely Continious Operators*. Springer Berlin, 1960.
- [32] W. Sun, *G-frames and G-Riesz bases*, *Journal of Mathematical Analysis and Appl*, 322, Issue. 1, pp. 437–452, 2006.
- [33] K. Zhuo, *Operator Theory in Function Spaces*, Marcel Dekker, Inc, 1990.